



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

heteronomous tarsi. On my last visit to Washington Col. Casey kindly showed me the type in which I recognized a Clerid and possibly the same as *Thaneroclerus tantillus* which species it seemed to resemble very closely. Unfortunately that species was not represented in Col. Casey's collection though no comparison could be made. However, the next day Mr. Schwarz showed me specimens of that species in the Nat. Museum and judging by memory I did not see any reason to change my opinion expressed the day before to Col. Casey, except that his specimen was perhaps a little more finely punctured than those in the National Museum collection. As I do not possess a specimen of *tantillus* nor *crinita* I am not able to state definitely that it is that species. Wolcott, erecting the genus *Prionodera* for *T. tantillus* describes the tarsi as four-jointed while Casey states that they are heteronomous in *Ababa crinita*. The first joint of the front and middle tarsi are often very small and more or less covered by the second joint and therefore easily overlooked, especially when the joints are densely pubescent.

---

### MISCELLANEOUS NOTES.

#### **Records of *Cænomyia ferruginea* in the Vicinity of New York City.**

—This widely distributed fly is not recorded in the last list of New Jersey insects, so the following records may be of interest. West Point, N. Y., June 3 and 4, 1911, 10 males, 9 females; June 13, 1914, one male. Fort Montgomery, N. Y., May 31, 1903, female. Ramapo, N. Y., May 31, female; June 3, 1905, female. Greenwood Lake, N. J., June 6, 1909, male. Alpine, N. J., May 3, 1909, female. The above mentioned are in the author's collection. In the American Museum there are three specimens from the following localities: Suffern, N. Y., male. Pearl River, N. Y., May 31, 1913, female (F. E. Watson). Hewitt, N. J., June 8, 1913, male.

It will be noted that all of these localities are in the hill country. So far the insect has not been collected in the more level portions along the coast, such as Long Island, Staten Island and the plains of New Jersey. A box containing these flies has a peculiar and not unpleasant odor, resembling that of the bark or dried leaves of the slippery elm.—WM. T. DAVIS.

**Some Records of Bees.**—Some time ago I received a number of bees from the Queensland Museum, without any information as to collectors. Dr. J. Bequaert, to whom I mentioned the matter, at once recognized from the localities the source of the collection. It was sent out by E. Le Moults of Paris, who is at present advertising insects for sale in American entomological journals. Dr. Bequaert showed me a series of similar insects, from the same source, in his possession. As many collectors doubtless possess materials from Le Moults, and as the locality records are new, it may be worth while to record the following. We are still without information as to the actual collectors.

1. Guyand, Maroni. *Acanthopus splendidus* Fabr., *Aglæ cærulea* Lep., *Exarete frontalis* Guér., *Oxæa festiva* Smith, *Bombus incarum* Franklin, *Centris obsoleta* Lep., *C. americana* Klug, *Epicharis conica* Smith, *E. schrottkyi* Friese, *E. affinis* Smith, *Ceratina lata* Spinola, *Eulæma dimidata* L., *E. fasciata* Lep., *E. Mocsaryi* Friese, *E. smaragdina mexicana* Mocs., *Euglossa piliventris* Guér.

2. Tunis. *Melecta luctuosa meridionalis* Gribodo, *Osmia ferruginea* Latr.

3. Ras-el-ma, Algeria. *Andrena albopunctata* Rossi, *Anthidium siculum* Spinola, *A. manicatum barbarum* Lep.

4. Dimbroko, French W. Africa. *Crocisa excisa* Friese.

5. N. Djole, Gabon. *Megachile tricolor* Friese.

For other species see Ent. News, 1916, p. 461, and Can. Ent., 1916, p. 406.—T. D. A. COCKERELL.

---

## PROCEEDINGS OF THE NEW YORK ENTOMOLOGICAL SOCIETY.

### MEETING OF FEBRUARY 6.

A regular meeting of the New York Entomological Society was held at 8:15 P. M., February 6, 1917, in the American Museum of Natural History. In the absence of the President, Mr. George W. J. Angell acted as chairman, with 21 members and one visitor present.

Prof. Edwin E. Calder, 4 Market Sq., Providence, R. I., was elected an active member.

Mr. Bird read a paper, "A New *Papaipema* from the Prairies," illustrated